

Effect of Probiotics and Prebiotics on Oral Health

Goutam Nanavati¹ T. Prasanth¹ Manab Kosala¹ Sujit K. Bhandari² Pamil Banotra¹

¹Department of Periodontology, Armed Forces Medical College, Pune, Maharashtra, India

²Department of Oral and Maxillofacial Surgery, Armed Forces Medical College, Pune, Maharashtra, India

Address for correspondence T. Prasanth, MDS, Department of Periodontology, Armed Forces Medical College, Wanowrie, Pune 411040, Maharashtra, India (e-mail: tprasanthavin@gmail.com).

Dent J Adv Stud:2021;9:1–6

Abstract

Probiotics are the living microorganisms that have been commonly used in the prevention of gastrointestinal disorders. In recent times, probiotic and prebiotics have been used to assess and develop a natural balance of the microbial flora in the respiratory tract and the oral cavity as an adjunct therapy. They are known to augment the existing microbial flora that is beneficial to the host. Prebiotics are nondigestible food ingredients that help in increasing populations of probiotic bacteria. Recent studies have shown that probiotics help in active reduction in gingivitis, dental caries, periodontitis, and halitosis by replacing the harmful oral species, by means of utilizing abundant health-associated oral microbial species. Additionally, the nitrate-reducing bacteria have shown promising effect in improving efficiency of probiotic strains to accentuate oral health benefits. Probiotics along with prebiotics effectively alter the host–microbial interface by achieving homeostasis in multifactorial diseases such as periodontal diseases and oral malodor. The aim of the review is to collate the existing information available on use of probiotics and prebiotics in oral cavity.

Keywords

- ▶ probiotics
- ▶ prebiotics
- ▶ periodontitis
- ▶ halitosis

Introduction

Probiotics are live nonpathogenic microorganisms administered to improve microbial balance the host.¹ Probiotics together with prebiotics confer synergistic effects. The term “prebiotics” was first introduced by Gibson and Roberfroid in 1995, where they used *Bifidobacterium* and *Lactobacillus* to deliver health-promoting benefits by modulation of human colon.² Use of microbiome therapy in homeostasis of oral cavity is a relatively new concept.

Since the conceptualization of probiotics, they have been described in many different ways. In recent times, use of probiotics is not restricted only to the gut and intestine, but also in oral cavity, female urogenital tract, and respiratory tract.³ Thus, probiotics have been defined as “live microorganisms that, when administered in adequate amounts, confer a health benefit on the host.”⁴

Prebiotics are the compounds in food that add to the growth and/or increase the activity of the probiotic organisms in the host. These prebiotics are basically fibrous compounds that are nondigestible. Thus, they pass through upper gastrointestinal (GI) tract undigested, acting as the substrate for the advantageous microorganisms, helping in their growth and enhanced biological activity.³ One need not take prebiotic for probiotic organisms to act, but together prebiotics make probiotic organisms more effective.

Periodontitis is a chronic inflammatory disease characterized by connective tissue destruction and alveolar bone resorption.⁵ Most of the therapies are directed toward arresting the disease progression, regaining the lost periodontal tissue, and balancing the host response to the virulence factor of the pathogenic organisms. Socransky in 1998 explained different bacterial complexes, related to periodontium in

published online
February 8, 2021

DOI <https://doi.org/10.1055/s-0040-1722523>
ISSN 2321-1482.

©2021. Bhojia Dental College and Hospital affiliated to Himachal Pradesh University.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Thieme Medical and Scientific Publishers Pvt. Ltd. A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

health and disease. Red and orange complex bacteria are the most common microorganisms associated with periodontal disease.⁶

Halitosis is the presence of foul and unpleasant odor originating from the oral cavity, primarily caused by *Fusobacterium nucleatum*, *Porphyromonas gingivalis*, *P. intermedia*, and *Treponema denticola*. It is due to the production of volatile sulfur compounds (VSCs).⁷

Dental caries is an infectious microbial disease resulting from production of organic acids due to the fermentation of carbohydrate. These acids cause demineralization of hard tissue of tooth, resulting in destruction and loss of tooth structure.⁸

All the above conditions have microbial plaque as a common etiological factor. Probiotics play an important role in modifying the local environment of oral cavity and reducing the dysbiosis. Thus, they help in maintaining homeostasis.⁹ Keeping in mind the growing resistance to the antibiotics and frequent recolonization of the organisms in dental plaque, use of probiotics in oral cavity is on the rise.¹⁰

History of Probiotics

Probiotics were introduced as early as 1908 by Elie Metchnikoff where he demonstrated the health benefits and increased lifespan in Bulgarian population attributing to consumption of fermented dairy products.¹ Being the pioneer of given field, he is referred to as “grand father of modern probiotics.” Dairy products such as milk, curd, and buttermilk are a regular part of diet in India. Medieval literature on dietetics “Kshemakutuhalam” emphasizes the importance of including components of milk products in diet, calling buttermilk as “elixir of gods.”¹¹ Use of sour milk as a cure for upset stomach has been documented in Bible also.

Around early 1900s, Henry Tissier, a French pediatrician, also conducted a study on probiotic organisms. He observed that children suffering from diarrhea when ingested with Y shaped “bifid” bacteria presented with healthy gut flora.¹² “Yakult,” a commercially available probiotic drink consisting of “*Lactobacillus*” species, was launched in Japan in 1930.¹³ The term “probiotics” became popular after Lilly and Stillwell used it for the first time in 1965 to address health

effect of many protozoal species.¹⁴ *Lactobacillus acidophilus* (Hull et al) and *Bifidobacterium bifidum* (Holcomb et al) were the first probiotic microorganisms used for research purpose.^{15,16}

Haukioja et al are considered as early pioneers in studying the interaction of probiotics in oral cavity. They analyzed the adhesion of *lactobacilli* and *bifidobacterial* species to hydroxyapatite crystals that were coated with saliva (in vitro).¹⁷

Probiotics

Probiotics is a term derived from Greek language, which means “for life.” They include many microbial species and strains.¹⁸ Commonly studied probiotics applied in the food and health areas are given in ►Table 1.

Several studies (both clinical and experimental) have proven that certain GI tract bacteria, that is, *Lactobacillus* and *Bifidobacterium* species, can control and manipulate the growth of microorganisms causing oral diseases. For probiotics to confer its benefits to host, the food should contain at least 10⁶–10⁷ live organisms per g or mL of product at the time of consumption, general frequency of consumption of probiotics being five billion to 10 billion colony-forming units per day.¹⁹

Guidelines for Use of Probiotics

FAO/WHO in 2002 issued guidelines for the evaluation of probiotics for use in food products.

1. Probiotic organisms must be a live organism.
2. The organism must be identified up to species level.
3. It should have a proven safety data.
4. When used in a defined value and delivery system, must have a physiological effect on the host.
5. When used as food additive, must be biologically and genetically stable.
6. It should be economical.
7. It should maintain long-term stability on storage (adequate shelf life).
8. It should be nonpathogenic and nonhazardous.
9. It should be able to influence local metabolic activity.²⁰

Table 1 Commonly used probiotics

Species	Strains	References
Lactobacillus	<i>L. acidophilus</i> , <i>L. amylovorus</i> , <i>L. casei</i> , <i>L. cebollobiosus</i> , <i>L. crispatus</i> , <i>L. curvatus</i> , <i>L. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. fermentum</i> , <i>L. gasser</i> , <i>L. helveticus</i> , <i>L. johnsonii</i> , <i>L. lactis</i> , <i>L. paracasei</i> , <i>L. pentosus</i> , <i>L. plantarum</i> , <i>L. reuteri</i> , <i>L. rhamnosus</i> , <i>L. salivarius</i>	25,52–55
Bifidobacterium	<i>B. adolescentis</i> , <i>B. animalis</i> , <i>B. bifidum</i> , <i>B. breve</i> , <i>B. infantis</i> , <i>B. lactis</i> , <i>B. longum</i> , <i>B. thermophilus</i>	25,53,54
Other species of lactic acid bacteria	<i>Enterococcus faecium</i> , <i>Lactococcus lactis</i> , <i>L. lactis</i> subsp. <i>cremoris</i> , <i>Leuconostoc mesenteroides</i> , <i>Pediococcus acidilactici</i> , <i>Streptococcus salivarius</i> subsp. <i>thermophilus</i> , <i>Streptococcus thermophilus</i> , <i>S. intermedius</i> , <i>S. diaacetylactis</i> , <i>Sporolactobacillus inulinus</i>	53,54
Nonlactic acid bacteria	<i>Bacillus cereus</i> var. <i>toyoi</i> , <i>Bacillus coagulans</i> , <i>Bacillus clausii</i> , <i>Saccharomyces cerevisiae</i> , <i>Saccharomyces boulardii</i> , <i>Escherichia coli</i> strain nissle, <i>Propionibacterium freudenreichii</i>	54–56

Mechanism of Action of Probiotics

In recent times, probiotics have emerged as substitute to antibiotics to help reduce various oral conditions such as periodontitis, dental caries, and halitosis. The proposed mechanism of action can be broadly classified under direct and indirect. Direct mode of action comprises of probiotic organisms having effect on the pathogenic organisms itself (► Fig. 1). The indirect mode of action is the one with probiotic organisms modulating the host response toward the pathogens.

Direct Action of Probiotics

L. reuteri has bactericidal action by producing ethanol, reuterin, and reutericyclin. It also helps in production of cobalamin and folate in the gut of host (► Fig. 1).

L. jensenii, *L. johnsonii*, and *B. bifidum* damage the bacterial cell walls by releasing hydrogen peroxide. Many other probiotics produce bacteriocin enzyme, which destroys the pathogen. At the same time, the enzymatic biosurfactants secreted by the probiotic organisms help reduce adhesion of the bacteria to each other as well as to the plaque or/and tooth surface.²¹⁻²⁴

Indirect Action of Probiotics

- Modulation of systemic immune response in favor of the host
- Affect local immune response
- Production of antioxidants or act as an antioxidant
- Reduction of pathogen associated proinflammatory cytokine production
- Prevention of plaque formation by neutralization of free electron
- Upregulation of epithelial barrier integrity and mucous production²¹⁻²³

Prebiotics

The term was coined by Gibson and Roberfoid in 1995, where they defined prebiotics as “the substrate that is selectively utilized by host microorganisms conferring a health benefit.”²⁵ They are the food ingredients that are relatively not digested by host digestive system, which stimulate growth and activity of the probiotic organisms.

The most commonly used prebiotics are insulin type fructans, fructo-oligosaccharide, galacto-oligosaccharide, and lactose.²⁶

Characteristic features of ideal prebiotic:

- They are neither hydrolyzed nor absorbed by mammalian enzymes or tissue.
- They selectively promote growth of certain probiotic species.
- They should be able to alter intestinal microbial flora and its activity.
- They should be able to modify systemic aspects of host defense system.²⁷

Symbiosis

Symbiosis is a mutually beneficial relationship between different groups. In this scenario, the term is used for the commercially available product containing both probiotic and prebiotic component. Due to this combination, the survival rate of the probiotic organism increases, thus enhancing the chances of effective action of probiotic bacteria,²⁸ for example, Bifido bacterial species in conjugation with fructo-oligosaccharide and *Lactobacillus* species in conjugation with lactulose. This symbiosis helps improve the survivability and boost the growth of these probiotics.²⁷

Probiotics in Oral Health

The use of probiotics in enhancing oral health dates back to early 90s.

Probiotics and Periodontal Disease

Probiotic organisms act as a supportive tool with binary benefits to manage periodontal inflammation. Primarily by causing dysbiosis, they result in competitive inhibition of periodontal pathogen thereby reducing the microbial load.²⁹

Secondarily by invigorating dendritic cells, they cause increase in the number helper T cells that modulate host immune response toward microorganisms present with the added benefit of least periodontal destruction during the process.³⁰ Bacteriocins secreted by *L salivarius*, *L gasseri*, and catalase secreted by *Weissella cibaria* cause an alteration in aggregation of bacteria responsible for plaque formation.³⁰ Streptococcal strains when used as probiotics, as




Fig. 1 Direct action of probiotics in oral cavity.

an adjunct to scaling, and root planning in chronic periodontitis patient show a significant reduction in plaque score along with reduction in red complex bacteria.³¹ Reutrin and reutericyclin enzymes secreted by *L. reuteri* inhibit the growth of pathogens by preventing the secretion of proinflammatory mediators. As a result, they exert an anti-inflammatory effect.³² *L. gasseri* and *L. fermentum* are more prevalent in healthy periodontium as compared with patients with chronic periodontitis, due to the production of hydrogen peroxide and bacteriocin.³³ *Bifidobacterial* species have shown competitive inhibition of pigment forming bacteria in oral cavity because of its utilization of vitamin K as substrate.³⁴

It was observed, when *L. plantarum* (HKL-137) was used during supportive therapy for chronic periodontitis for 12 weeks; there was reduction in probing depth as compared with the placebo group.³⁵ *L. rhamnosus* and *B. lactis* (BB-12) affect the numbers of *P. gingivalis* and *A. actinomycetemcomitans* by significantly reducing their load in oral cavity and help in promoting gingival health.³⁶

Probiotics and Halitosis

Halitosis is the general term used to describe any disagreeable odor of expired air, regardless of its origin.³⁷ It is an unpleasant odor most commonly caused by *Fusobacterium nucleatum*, *P. gingivalis*, *P. intermedia*, and *T. denticola*. The microbiota causes putrefactive metabolism of amino acids producing VSCs (hydrogen sulfide, methyl mercaptan, dimethyl sulfide).³⁸

Tongue acts as a niche for accumulation of microbiota due to the presence of multiple papillae on its dorsal surface. Probiotics help to reduce the dysbiosis of microbiota on the tongue especially the posterior part, which is difficult to clean.³⁹

L. salivarius WB21 strain when administered orally as tablet, thrice a day, showed a significant decrease in organoleptic score due to decline in the concentration of VSCs.⁴⁰

W. cibaria when used as a probiotic agent demonstrated a reduction in H₂O₂ production, due to inhibition of *F. nucleatum*.⁴¹ *Streptococcus salivarius* (K12 strain), one of the pioneer probiotic strain, secretes bacteriocin, which when consumed as lozenges helps reduce *Solobacterium moorei*, thereby reducing oral mal odor.⁴²

L. reuteri has also been beneficial in promoting oral health and reducing oral mal odor. By downregulation of enzyme methionine γ lyase is produced by *F. nucleatum* and *P. gingivalis*.⁴³

Probiotics and Dental Caries

Caries is an infectious microbial disease of oral cavity caused by fermentation of carbohydrates by bacteria resulting in dissolution of the mineral content of the tooth structure. The main organism responsible is *S. mutans*. There is increased prevalence of dental caries, especially in children due to increase in consumption of refined sugar and inadequate cleaning habits.⁴⁴

L. rhamnosus GG strain, *L. reuteri*, and *Bifidobacterium lactis* BB-12 strain when used as probiotics have shown

significant reduction in the number of *S. mutans*. They potentially deter the microbiota of dental plaque and reduce the adhesion of bacteria to the tooth surface, resulting in reduced incidence of dental caries. When multispecies or multistrains of probiotic organisms are used, there is a considerable decline in the cariogenic bacteria.⁴⁵ Similarly, *L. rhamnosus* 1b21 strain, when used in milk along with fluoride, produces the same result.⁴⁶

Commercially Available Probiotics

Probiotics used to elevate oral health and treat dental conditions are commonly available as lozenges, toothpaste, chewing gums, or mouthwash.

ProBiora Health

It is designed to improve in general oral health of humans as well as pets. It is a multistrain blend of *Streptococcus* strains, *S. oralis* KJ3, *S. uberis* KJ2, and *S. rattus* JH145.⁴⁷

Gum PerioBalance

It is one of the pioneer formulations for treating periodontal disease. It has a combination of two strains of *L. reuteri* to act against periodontal pathogens.⁴⁸

Blis Technologies

It includes the formulation of two strains of BLIS K12 and BLIS M18 probiotics that are native to oral cavity. They control halitosis along with having beneficial effect on throat and nasal health.⁴⁹

PeriBiotic

It is a toothpaste containing *L. paracasei* probiotic that reduces periodontal pathogens.⁴⁸

NatureWise

This probiotic preparation is commercially available in the form of chewable tablets. It contains 12 different types of probiotic strains: *L. casei* subsp. *casei*, *L. paracasei* Lpc-37, *L. plantarum*, *L. reuteri*, *L. salivarius*, *S. salivarius* K12, *S. salivarius* M18, *B. lactis* BI-04, *L. rhamnosus* GG, *B. breve* Bb-03, *B. infantis*, and *S. thermophilus*. The aim is to have a protective effect on the oral cavity by altering the pathogenic bacterial colonies.

Prodentis

This probiotic lozenge is a blend of two *Lactobacillus reuteri* strains containing a minimum of 1×10^8 CFU for each of the strains DSM 17938 and ATCC PTA 5289.⁴⁸

Safety Concerns

FDA classifies probiotic organisms as generally regarded as safe. But due to the increase in probiotic supplementation of various food products in the recent years, their safety standards still need to be investigated and verified. There is still a reasonable likelihood of bacteremia in critically ill or immunocompromised patients as probiotics are live organisms.⁵⁰ It

may also present as a potential risk for people with lactose intolerance or hypersensitivity to milk products.

Future of Probiotics

With an increasing threat of microorganisms developing resistance to the known antibiotics, probiotics have emerged as an improved and innovative solution to the above-stated problem. Probiotics are way ahead of their conventional counter products with an added advantage of minimal side effects. There is a change in lifestyle and feeding habits of people across the globe, indicating an upsurge of oral diseases proportional to other lifestyle diseases attributing to the same factor. These natural strains of probiotic organisms isolated from dairy products, for example, curd, buttermilk can be used on regular basis for management and maintenance of dental diseases and oral health, respectively.

“Designer probiotics,” a relatively new concept of genetically modified microorganisms which target a specific pathogen, can be utilized as a novel tool to act against the severe oral diseases. These strains achieve the desired effect by disruption of biofilm and formation of enzymes such as bacteriocins. They can also be designed to endure high amount of stress, thus improving their sustainability in oral cavity. Successful use of probiotics in dentistry can pave a way to advanced treatment options and therapy modules to augment the oral health in general.⁵¹

Conclusion

The aim of this review was to find out the existing literature regarding the use of probiotics in oral cavity. Probiotics as we know are useful in maintaining the microbial balance in the human body. Use of probiotics in oral cavity is an emerging concept. They play a significant role in clinical management of halitosis, dental caries, and periodontal diseases. These studies suggest that probiotics can reduce the load of pathogenic organisms as well as restore the host microbiota. To explore the potential of prebiotics, further studies are required to understand the ability of the probiotic organism to survive, grow, and have a therapeutic effect. Also we need well designed, long-term follow-up studies to fix the dosages and schedules of administration, interaction with various substrate, interpretation of oral health risks in case of long-term use of the probiotic organisms; so as to emphasize and practice the same accordingly.

Conflict of Interest

None declared.

References

- 1 Metchnikoff E, *Optimistic Studies*. New York: Putman's Sons; 1908 ;161–183
- 2 Gibson GR, Roberfroid MB. Dietary modulation of the human colonic microbiota: introducing the concept of prebiotics. *J Nutr* 1995;125(6):1401–1412
- 3 Gibson GR, Hutkins R, Sanders ME, et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol* 2017;14(8):491–502
- 4 Food and Agriculture Organization and World Health Organization Expert Consultation, Evaluation of health and nutritional properties of powder milk and live lactic acid bacteria. Córdoba, Argentina: Food and Agriculture Organization of the United Nations and World Health Organization; 2001
- 5 Page RC, Kornman KS. The pathogenesis of human periodontitis: an introduction. *Periodontol* 2000 1997;14(1):9–11
- 6 Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL Jr. Microbial complexes in subgingival plaque. *J Clin Periodontol* 1998;25(2):134–144
- 7 Persson S, Edlund MB, Claesson R, Carlsson J. The formation of hydrogen sulfide and methyl mercaptan by oral bacteria. *Oral Microbiol Immunol* 1990;5(4):195–201
- 8 Featherstone JD. The continuum of dental caries—evidence for a dynamic disease process. *J Dent Res* 2004;83 Spec No C(1–suppl) :C39–C42
- 9 Allaker RP, Stephen AS. Use of probiotics and oral health. *Curr Oral Health Rep* 2017;4(4):309–318
- 10 Zambori C, Morvay AA, Sala C, et al. Antimicrobial effect of probiotics on bacterial species from dental plaque. *J Infect Dev Ctries* 2016;10(3):214–221
- 11 Kukkupuni SK, Shashikumar A, Venkatasubramanian P. Fermented milk products: probiotics of ayurveda. *Journal of Medical Nutrition and Nutraceuticals*. 2015;4(1):14–21
- 12 Tissier H. The treatment of intestinal infections by the method of transformation of bacterial intestinal flora. *CR Soc. Biol.* 1906;60:359–361
- 13 Anukam KC, Reid G. Probiotics: 100 years (1907–2007) after Elie Metchnikoff's observation. *Communicating current research and educational topics and trends in applied microbiology*; 2007;1:466–474
- 14 Lilly DM, Stillwell RH. Probiotics: growth-promoting factors produced by microorganisms. *Science* 1965;147(3659):747–748
- 15 Hull RR, Roberts AV, Mayes JJ. Survival of *Lactobacillus acidophilus* in yoghurt. *Australian Journal of Dairy Technology* 1984;39(4):164
- 16 Holcomb JE, Frank JF, Mc Gregor JU. Viability of *Lactobacillus acidophilus* and *Bifidobacterium bifidum* in softserve frozen yogurt. 1991;26:4–5
- 17 Haukioja A, Yli-Knuuttila H, Loimaranta V, et al. Oral adhesion and survival of probiotic and other lactobacilli and bifidobacteria in vitro. *Oral Microbiol Immunol* 2006;21(5):326–332
- 18 Isolauri E. Probiotics in human disease. *Am J Clin Nutr* 2001;73(6):1142S–1146S
- 19 Shah NP. Functional cultures and health benefits. *Int Dairy J* 2007;17(11):1262–1277
- 20 Joint FAO/WHO Expert Committee on Food Additives, Joint FAO/WHO Expert Committee on Food Additives. Meeting, World Health Organization. Evaluation of Certain Food Additives and Contaminants: Fifty-seventh Report of the Joint FAO/WHO Expert Committee on Food Additives. World Health Organization; 2002 Dec 13
- 21 Mahasneh SA, Mahasneh AM. Probiotics: a promising role in dental health. *Dent J (Basel)* 2017;5(4):26–36
- 22 Agarwal G, Ingle NA, Kaur N, Yadav P, Ingle E, Charania Z. Probiotics and oral health: a review. *J Int Oral Health* 2015;7(10):133–136
- 23 Meurman JH. Probiotics: do they have a role in oral medicine and dentistry? *Eur J Oral Sci* 2005;113(3):188–196
- 24 Haukioja A. Probiotics and oral health. *Eur J Dent* 2010;4(3):348–355
- 25 Bustamante M, Oomah BD, Rubilar M, Shene C. Effective *Lactobacillus plantarum* and *Bifidobacterium infantis* encapsulation with chia seed (*Salvia hispanica* L.) and flaxseed

- (*Linum usitatissimum* L.) mucilage and soluble protein by spray drying. *Food Chem* 2017;216:97–105
- 26 Pandey KR, Naik SR, Vakil BV. Probiotics, prebiotics and synbiotics- a review. *J Food Sci Technol* 2015;52(12):7577–7587
 - 27 Maftai NM. Probiotic, Prebiotic and Synbiotic Products in Human Health. In: *Frontiers and New Trends in the Science of Fermented Food and Beverages*. London, IntechOpen;2019
 - 28 Sekhon BS, Jairath S. Prebiotics, probiotics and synbiotics: an overview. *Journal of Pharmaceutical Education and Research*. 2010;1(2):13–36
 - 29 Saha S, Tomaro-Duchesneau C, Tabrizian M, Prakash S. Probiotics as oral health biotherapeutics. *Expert Opin Biol Ther* 2012;12(9):1207–1220
 - 30 Bose BB. Probiotics-promotes periodontal health?-An insight. *International Journal of Dental Sciences and Research*. 2013;1(3):67–70
 - 31 Laleman I, Yilmaz E, Ozelik O, et al. The effect of a streptococci containing probiotic in periodontal therapy: a randomized controlled trial. *J Clin Periodontol* 2015;42(11):1032–1041
 - 32 Szkaradkiewicz AK, Stopa J, Karpiński TM. Effect of oral administration involving a probiotic strain of *Lactobacillus reuteri* on pro-inflammatory cytokine response in patients with chronic periodontitis. *Arch Immunol Ther Exp (Warsz)* 2014;62(6):495–500
 - 33 Gupta G. Probiotics and periodontal health. *J Med Life* 2011; 4(4):387–394
 - 34 Hojo K, Nagaoka S, Murata S, Taketomo N, Ohshima T, Maeda N. Reduction of vitamin K concentration by salivary *Bifidobacterium* strains and their possible nutritional competition with *Porphyromonas gingivalis*. *J Appl Microbiol* 2007;103(5):1969–1974
 - 35 Iwasaki K, Maeda K, Hidaka K, Nemoto K, Hirose Y, Deguchi S. Daily intake of heat-killed *Lactobacillus plantarum* L-137 decreases the probing depth in patients undergoing supportive periodontal therapy. *Oral Health Prev Dent* 2016;14(3):207–214
 - 36 Alanzi A, Honkala S, Honkala E, Varghese A, Tolvanen M, Söderling E. Effect of *Lactobacillus rhamnosus* and *Bifidobacterium lactis* on gingival health, dental plaque, and periodontopathogens in adolescents: a randomised placebo-controlled clinical trial. *Benef Microbes* 2018;9(4):593–602
 - 37 Aydin M, Harvey-Woodworth CN. Halitosis: a new definition and classification. *Br Dent J* 2014;217(1):E1–E10
 - 38 Niles HP. *Advances in Mouth Odor Research. Bad Breath: A Multidisciplinary Approach*; 1996:55–69
 - 39 Gurpinar B, Yildirim G, Kumral TL, et al. A simple method to reduce halitosis; tongue scraping with probiotics. *J Breath Res* 2019;14(1):016008
 - 40 Suzuki N, Yoneda M, Takeshita T, Hirofuji T, Hanioka T. Induction and inhibition of oral malodor. *Mol Oral Microbiol* 2019;34(3):85–96
 - 41 Kang MS, Kim BG, Chung J, Lee HC, Oh JS. Inhibitory effect of *Weissella cibaria* isolates on the production of volatile sulphur compounds. *J Clin Periodontol* 2006;33(3):226–232
 - 42 Wescombe PA, Heng NC, Burton JP, Tagg JR. Something old and something new: an update on the amazing repertoire of bacteriocins produced by *Streptococcus salivarius*. *Probiotics Antimicrob Proteins* 2010;2(1):37–45
 - 43 Fujiwara N, Murakami K, Nakao M, et al. Novel reuterin-related compounds suppress odour by periodontopathic bacteria. *Oral Dis* 2017;23(4):492–497
 - 44 Tinanoff N, Baez RJ, Diaz Guillory C, et al. Early childhood caries epidemiology, aetiology, risk assessment, societal burden, management, education, and policy: Global perspective. *International journal of Paediatric Dentistry* 2019;29(3):238–248
 - 45 Meurman JH, Stamatova I. Probiotics: contributions to oral health. *Oral Dis* 2007;13(5):443–451
 - 46 Galdeano CM, de Moreno de LeBlanc A, Vinderola G, Bonet ME, Perdígón G. Proposed model: mechanisms of immunomodulation induced by probiotic bacteria. *Clin Vaccine Immunol* 2007;14(5):485–492
 - 47 Duarte C, Al-Yagoob A, Al-Ani A. Efficacy of probiotics used as a periodontal treatment aid: a pilot study. *Saudi Dent J* 2019;31(1):143–147
 - 48 Lawande S. Probiotics for management of periodontal disease: a novel therapeutic strategy. *IOSR J Pharm* 2012;2:41–46
 - 49 Zupancic K, Kriksic V, Kovacevic I, Kovacevic D. Influence of oral probiotic *Streptococcus salivarius* K12 on ear and oral cavity health in humans: systematic review. *Probiotics Antimicrob Proteins* 2017;9(2):102–110
 - 50 Cannon JP, Lee TA, Bolanos JT, Danziger LH. Pathogenic relevance of *Lactobacillus*: a retrospective review of over 200 cases. *Eur J Clin Microbiol Infect Dis* 2005;24(1):31–40
 - 51 Chatterjee A, Bhattacharya H, Kandwal A. Probiotics in periodontal health and disease. *J Indian Soc Periodontol* 2011;15(1):23–28
 - 52 Collins MD, Gibson GR. Probiotics, prebiotics, and synbiotics: approaches for modulating the microbial ecology of the gut. *Am J Clin Nutr* 1999;69(5):1052S–1057S
 - 53 Holzapfel WH, Haberer P, Geisen R, Björkroth J, Schillinger U. Taxonomy and important features of probiotic microorganisms in food and nutrition. *Am J Clin Nutr* 2001; 73(2, Suppl):365S–373S
 - 54 Markowiak P, Śliżewska K. Effects of probiotics, prebiotics, and synbiotics on human health. *Nutrients* 2017;9(9):1021–1051
 - 55 Bustamante M, Oomah BD, Mosi-Roa Y, Rubilar M, Burgos-Díaz C. Probiotics as an adjunct therapy for the treatment of halitosis, dental caries and periodontitis. *Probiotics Antimicrob Proteins* 2020;12(2):325–334
 - 56 Endres JR, Clewell A, Jade KA, Farber T, Hauswirth J, Schauss AG. Safety assessment of a proprietary preparation of a novel Probiotic, *Bacillus coagulans*, as a food ingredient. *Food Chem Toxicol* 2009;47(6):1231–1238